


Bioinspired Total Synthesis of Bussealin E

David G. Twigg, Leonardo Baldassarre, Elizabeth C. Frye, Warren R. J. D. Galloway, and David R. Spring*¹

Department of Chemistry, University of Cambridge, Lensfield Road, Cambridge CB2 1EW, U.K.

S Supporting Information



ABSTRACT: The first total synthesis of bussealin E, a natural product with a unique cycloheptadibenzofuran scaffold, is reported. A strategy inspired by a proposed biosynthesis was employed whereby a diphenylpropane derivative underwent an oxidative phenolic coupling to forge the tetracyclic ring system. The synthesis of the diphenylpropane featured a key sp^2 – sp^3 Hiyama coupling between a vinyldisiloxane and a benzylic bromide.

First isolated in 2007 from the roots of the endemic Malagasy plant *Bussea sakalava*, the bussealin natural products consist of four diphenylpropanes (A–D) and a cycloheptadibenzofuran (E, **1**, Figure 1) which display




Figure 1. Bussealin natural products.


moderate antiproliferative activity against the A2780 human ovarian cancer cell line (IC_{50} 24–45 μM).¹ More thorough biological assessment has been hampered by insufficient material, and expedient synthetic routes are required to investigate this natural product family further. In light of this, the first total synthesis of bussealin A was reported in 2012, with biological profiling indicating similar low micromolar levels of activity against two other cancer cell lines.² Of the other family members, bussealin E is notable for its unique carbon skeleton, with no other reported natural product bearing the same tetracyclic core. This structural novelty, alongside the potential biological applications of this unexplored system, suggested bussealin E as a target for total synthesis.

Noting that bussealin E was found alongside several diphenylpropanes, Pan et al. proposed a biosynthetic route featuring the oxidative cyclization and subsequent dehydration of an appropriately substituted intermediate **2** (Scheme 1A).¹

With such a biomimetic approach deemed attractive, our retrosynthetic analysis focused on diphenylpropane species **2**.

Using disconnections analogous to those employed in the synthesis of bussealin A,² it was anticipated that **2** could be generated from the corresponding protected olefin **3**, which we

Scheme 1. Proposed Biosynthesis (A) and Retrosynthesis (B) of Bussealin E (**1**)




Received: January 30, 2018

Published: March 2, 2018

envisaged accessing via an sp^2 – sp^3 Hiyama coupling of disiloxane **4** with benzyl bromide **5**, a reliable and atom-economical methodology previously developed in these laboratories (Scheme 1B).^{2,3} It was anticipated that the sequence of bromination, Sonogashira coupling, and hydrosilylation would deliver the disiloxane **4** from symmetrical tetraether **6**, itself accessible in multigram quantities from readily available 2,3,4-trimethoxybenzaldehyde **7** using established protocols. The synthesis of bromide **5** from methyl gallate **8** has been described previously.²

Synthetic efforts began with a Baeyer–Villiger oxidation of aldehyde **7**, followed by hydrolysis of the crude formate ester. Subsequent oxidation of the intermediate phenol using ceric ammonium nitrate on silica provided 2,3-dimethoxy-1,4-benzoquinone **9** (Scheme 2). Reduction with NaBH_4 and bis-

Scheme 2. Synthesis of Alkyne Intermediate 12




protection of the resultant hydroquinone with benzyl bromide proceeded smoothly to furnish tetraether intermediate **6**. Bromination was achieved in a near-quantitative yield using the conditions of Krishna Mohan et al. (Oxone, NH_4Br)⁴ to yield **10**, which then underwent Sonogashira cross-coupling with trimethylsilylacetylene and subsequent deprotection of the silyl group to give the desired terminal alkyne **12** in a combined 63% yield. The Sonogashira coupling required relatively high catalyst loadings (10 mol % $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$ and CuI) and an elevated temperature, possibly due to the electron-rich nature of the aryl bromide **10**. Intermediate purification of the trimethylsilyl-protected alkyne **11** was also found to improve the overall yield compared to a one-pot procedure which only gave the desired product **12** in 44% yield from bromide **10**.

Pt-catalyzed hydrosilylation of alkyne **12**, as per the conditions of Denmark et al.,⁵ generated the desired (*E,E*) isomer of disiloxane **4** in excellent yield and geometric purity (Scheme 3). Pleasingly, the key sp^2 – sp^3 Hiyama coupling of disiloxane **4** and benzyl bromide **5** (prepared in four steps from readily available methyl gallate **8**, see ref 2) proceeded in 80% yield and with complete retention of the olefin stereochemistry, with no evidence of the α isomer occasionally formed in such reactions.⁶ Catalytic hydrogenation of **3** over palladium on carbon then served to reduce the olefin and remove the four benzyl ether protecting groups, generating the desired tetraphenol species **2**.


With **2** in hand, investigation of the proposed oxidative phenolic coupling toward bussealin E commenced. A range of

Scheme 3. Synthesis of Key Diphenylpropane 2



oxidants have previously been used to effect similar intramolecular cyclizations.⁷ However, no examples featuring a 1,4-hydroquinone have been reported. Nevertheless, diphenylpropane **2** was subjected to a range of conditions, including metal salts such as MoCl_5 ,⁸ SnCl_4 ,⁹ VOF_3 ,¹⁰ $\text{Pb}(\text{OAc})_4$,¹¹ FeCl_3 ,¹² and the TMEDA complex of $\text{CuCl}(\text{OH})$;¹³ hypervalent iodine species $\text{PhI}(\text{OAc})_2$ and $\text{PhI}(\text{O}_2\text{CCF}_3)_2$;¹⁴ and the organic oxidant, 2,3-dichloro-5,6-dicyano-*p*-benzoquinone (DDQ).¹⁵ In all cases, a complex mixture of products was obtained, with the predominant species consistently observed by NMR analysis to be acyclic benzoquinone **13** (arising from oxidation of the corresponding 1,4-hydroquinone moiety of **2**) and the spirocyclic species **14**, which exists as a 3:1 mixture in solution with its hemiacetal isomer **15** (Scheme 4). Desired product **1**

Scheme 4. Synthesis and Proposed Mechanism of Formation of Bussealin E (1)



was only observed in trace amounts in some of these reactions, as assessed by comparing the analytical HPLC traces of the crude reaction mixtures against a reference sample of natural bussealin E (kindly provided for analysis by Prof. D. Kingston, Virginia Polytechnic Institute and State University). Of the reagents evaluated, FeCl_3 gave the most promising result in this regard; further exploration of this oxidant was carried out by varying temperature and reagent stoichiometry and including or excluding an acidic additive. Gratifyingly, the use of FeCl_3 adsorbed onto silica gel, as employed by Jempty et al.,¹⁶

followed by heating with *para*-toluenesulfonic acid produced a sufficient amount of bussealin E **1** to isolate an analytically pure sample, albeit in a low yield. While further attempts to improve this result were not successful, the ^1H and ^{13}C NMR spectra of the synthetic material closely matched those previously reported for the natural sample (see the *SI*), confirming its identity.

The concomitant observation of the previously mentioned byproducts **13** and **14** provides new insights into the possible biosynthesis of bussealin E (**1**). These results suggest a mechanism for the formation of **1** from **2** that proceeds via intramolecular Michael addition of initially generated benzoquinone **13**, with formation of the smaller 6-membered spirocyclic ring (path b, *Scheme 4*) favored over that of the 7-membered carbocycle (path a, *Scheme 4*), as evidenced by the reactions in this report. One previous publication demonstrates a similar result for an analogous diphenylethane system.¹⁷ When heated strongly in an acidic solution, an isolated sample of compound **14** did not undergo any change, with no observed rearrangement or retro-Michael reaction, suggesting **14** exists as a thermodynamically stable species. Alongside this observation, the low isolated yield of **1** may suggest that the reaction relies on an enzyme-catalyzed process when formed naturally in order to generate appreciable quantities of the natural product.

In conclusion, we have completed the first reported synthesis of the unique cycloheptadibenzofuran natural product, bussealin E. The final intermediate in the synthesis, diphenylpropane **2**, was prepared in 11 steps, in its longest linear sequence, in an overall 14% yield, featuring a key $\text{sp}^2\text{-sp}^3$ Hiyama coupling between a vinylsiloxane and a benzylic bromide. The final bioinspired cyclization to deliver bussealin E only proceeded in low yield, with the observed byproducts suggesting a mechanism that proceeds via Michael addition of the initially formed 1,4-benzoquinone **13** with selectivity for the undesired 6-membered ring as seen in compound **14**. This low yield could suggest that the reaction relies on an enzyme-catalyzed process when formed naturally. With a route to bussealin E determined, further synthetic studies and biological evaluation of this novel scaffold are underway and will be reported in due course.

■ ASSOCIATED CONTENT

■ Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: [10.1021/acs.orglett.8b00340](https://doi.org/10.1021/acs.orglett.8b00340).

Experimental details and characterization data for all products including ^1H and ^{13}C NMR spectra (*PDF*)

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: spring@ch.cam.ac.uk.

ORCID

David R. Spring: [0000-0001-7355-2824](https://orcid.org/0000-0001-7355-2824)

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This research was supported by the EPSRC, BBSRC, MRC, Wellcome Trust, and ERC (FP7/2007-2013; 279337/DOS). D.G.T. thanks AstraZeneca for funding. D.R.S. acknowledges support from a Royal Society Wolfson Research Merit award. We thank Prof. D. Kingston (Virginia Polytechnic Institute and State University) for providing a sample of authentic bussealin E. Data accessibility: all data supporting this study are provided as *Supporting Information* accompanying this paper.

■ REFERENCES

- (1) Pan, E.; Harinantenaina, L.; Brodie, P. J.; Miller, J. S.; Callmender, M. W.; Rakotonandrasana, S.; Rakotobe, E.; Rasamison, V. E.; Kingston, D. G. I. *J. Nat. Prod.* **2010**, *73*, 1792–1795.
- (2) Frye, E. C.; O' Connor, C. J.; Twigg, D. G.; Elbert, B.; Laraia, L.; Hulcoop, D. G.; Venkitaraman, A. R.; Spring, D. R. *Chem. - Eur. J.* **2012**, *18*, 8774–8779.
- (3) (a) Sore, H. F.; Boehner, C. M.; Laraia, L.; Logoteta, P.; Prestinari, C.; Scott, M.; Williams, K.; Galloway, W. R. J. D.; Spring, D. R. *Org. Biomol. Chem.* **2011**, *9*, 504–515. (b) Sore, H. F.; Boehner, C. M.; MacDonald, S. J. F.; Norton, D.; Fox, D. J.; Spring, D. R. *Org. Biomol. Chem.* **2009**, *7*, 1068–1071. (c) Boehner, C. M.; Frye, E. C.; O'Connell, K. M. G.; Galloway, W. R. J. D.; Sore, H. F.; Garcia-Dominguez, P.; Norton, D.; Hulcoop, D. G.; Owen, M.; Turner, G.; Crawford, C.; Horsley, H.; Spring, D. R. *Chem. - Eur. J.* **2011**, *17*, 13230–13239.
- (4) Krishna Mohan, K. V. V.; Narender, N. *Synthesis* **2012**, *44*, 15–26.
- (5) Denmark, S. E.; Wang, Z. *Org. Lett.* **2001**, *3*, 1073–1076.
- (6) (a) Yoshida, J.; Tamao, K.; Yamamoto, H.; Kakui, T.; Uchida, T.; Kumada, M. *Organometallics* **1982**, *1*, 542–549. (b) Denmark, S. E.; Choi, J. Y. *J. Am. Chem. Soc.* **1999**, *121*, 5821–5822.
- (7) Taylor, W. I.; Battersby, A. R. *Oxidative Coupling of Phenols*; M. Dekker, Inc.: New York, 1967.
- (8) Kramer, B.; Fröhlich, R.; Waldvogel, S. R. *Eur. J. Org. Chem.* **2003**, *2003*, 3549–3554.
- (9) Takeya, T.; Doi, H.; Ogata, T.; Otsuka, T.; Okamoto, I.; Kotani, E. *Tetrahedron* **2004**, *60*, 6295–6310.
- (10) Kupchan, S. M.; Dhingra, O. P.; Kim, C.-K. *J. Org. Chem.* **1978**, *43*, 4076–4081.
- (11) Feldman, K. S.; Ensel, S. M.; Minard, R. D. *J. Am. Chem. Soc.* **1994**, *116*, 1742–1745.
- (12) Bolm, C.; Legros, J.; Le Pailh, J.; Zani, L. *Chem. Rev.* **2004**, *104*, 6217–6254.
- (13) Ohishi, T.; Kojima, T.; Matsuoka, T.; Shiro, M.; Kotsuki, H. *Tetrahedron Lett.* **2001**, *42*, 2493–2496.
- (14) (a) Zhdankin, V. V.; Stang, P. J. *Chem. Rev.* **2002**, *102*, 2523–2584. (b) Pouységu, L.; Deffieux, D.; Quideau, S. *Tetrahedron* **2010**, *66*, 2235–2261.
- (15) Bohmdorfer, S.; Patel, A.; Gille, L.; Netscher, T.; Mereiter, K.; Rosenau, T. *Curr. Org. Synth.* **2013**, *10*, 165–168.
- (16) (a) Jemphy, T. C.; Miller, L. L.; Mazur, Y. *J. Org. Chem.* **1980**, *45*, 749–751. (b) Jemphy, T. C.; Gogins, K. A. Z.; Mazur, Y.; Miller, L. L. *J. Org. Chem.* **1981**, *46*, 4545–4551.
- (17) Thangaraj, S.; Tsao, W.-S.; Luo, Y.-W.; Lee, Y.-J.; Chang, C.-F.; Lin, C.-C.; Uang, B.-J.; Yu, C.-C.; Guh, J.-H.; Teng, C.-M. *Tetrahedron* **2011**, *67*, 6166–6172.